Insertion and Maintenance of Nasogastric Feeding Tubes in Infants, Children and Young People-Policy Toolkit- Appendix 1.

Appendix 1- Nasogastric Tube Position Confirmation & Recording Form

Addressograph or
Name:
DOB:
CRN:

It is mandatory to check pH:

- 1. following initial insertion
- 2. before administering each feed
- 3. before giving medication
- 4. at least twice daily during continuous feeds
- 5. following episodes of vomiting, retching, coughing, colour change
- 6. following evidence of tube displacement (for example, loose tape or visible tube appears longer)

Date/time tube inserted	Tube inserted by 1. Name 2. Job Title	Type of tube	Centimetre marking at nose or external tube length	pH on insertion	Date tube removed and reason

Date	Time	Checked by *Name *Job title	Any concerns re NG tube displacement	Centimetre marking at nose or external tube length	pH of Aspirate (if pH of aspirate is not within range refer to decision tree)	Safe to commence feed? Yes or No
_						
_						

Insertion and Maintenance of Nasogastric Feeding Tubes in Infants, Children and Young People-Policy Toolkit- Appendix 1.

NG Recording Form Continuation sheet

Addressograph or	
Name:	
DOB:	
CRN:	

Date	Time	Checked by *Name *Job title	Any concerns re NG tube displacement	Centimetre marking at nose or external tube length	pH of Aspirate (if pH of aspirate is not within range refer to decision tree)	Safe to commence feed? Yes or No
					Í	

ADILIDO	

Insertion and Maintenance of Nasogastric Feeding Tubes in Infants, Children and Young People-

Policy Toolkit- Appendix 1.